

Review

Inactivation of viral antigens for vaccine preparation with particular reference to the application of binary ethylenimine

Haas G. Bahnmann

Viral antigens for human and veterinary vaccines are still inactivated with formaldehyde. This is not an ideal inactivant and the problems of formaldehyde inactivation of vaccines are discussed. Vaccines inactivated with aziridines are superior in safety and antigenicity. Aziridines inactivate viruses in a first-order reaction and the inactivation rate and endpoint can be determined. The preparation and application of the aziridine compound binary ethylenimine (BEI) and the necessary conditions for and controls of the inactivation process are described and discussed. A computer program has been written for assistance in the use of BEI for controlled inactivation of viral antigens.

Keywords: Inactivation; binary ethylenimine; viral antigens

Introduction

A perusal of recent and current literature on the preparation of viral vaccines with inactivated antigen and, in particular, of experimental vaccines of this type shows that very often the inactivation is still obtained with formaldehyde and without the necessary controls. In the preparation of an inactivated viral vaccine the inactivation process is a very important step. The innocuity of the vaccine must be assured before the question of potency can be addressed.

It seems worthwhile to discuss the inactivation of viral antigens with reference to a few older (and perhaps forgotten) publications as well as some more recent studies. The procedures and process controls which must be applied in order to assure a safely inactivated vaccine will be described. The example of foot-and-mouth disease (FMD) vaccine preparation will be used as this vaccine is by volume the largest viral vaccine produced at present.

Inactivation with formaldehyde

For many years most of the viral vaccines with inactivated antigen were prepared with formaldehyde as inactivating agent. The work of Sven Gard and his collaborators^{1,2} with poliovirus during the period of 1956 to 1958 demonstrated that the inactivation of this virus with formaldehyde was not a linear or first-order reaction. Similar results were obtained for the formaldehyde inactivation of FMD virus by Wesslén and Dinter in 1957³ and by Graves⁴ in 1963.

A recent publication stating the linearity of formaldehyde inactivation of FMD virus⁵ comes to this erroneous conclusion because the infectivity titration for the inactivation slope was based on final readings of plaque forming units (p.f.u.) at 2 days. This is far too

short a time for a reading with formaldehyde treated virus which is unique in having a markedly extended incubation period for the first replication cycle in cell cultures, as was shown by Schultz *et al.* in 1957⁶ and Böttiger *et al.* in 1958⁷. For points on the lower part of the inactivation slope Böttiger *et al.* needed 12 days to obtain a final p.f.u. reading.

The extended incubation period for formaldehyde treated virus also means that an innocuity test in animals is inappropriate for detecting small amounts of residual infectious virus. When this virus begins to replicate with a delay of between several days and 2 weeks, the animal is already beginning to produce antibodies. The new virus will then be neutralized by antibody and the animal has an abortive or subclinical infection. This subclinical infection can be detected by testing the animal for the virus infection associated antigen (VIAA), the viral RNA polymerase⁸.

In 1975 Alonso *et al.*⁹ reported a study of cattle exposed to FMD and found that at 21 days postvaccination, of 18 animals vaccinated with formaldehyde inactivated vaccine five animals were positive for VIAA. Of 16 animals vaccinated with *N*-acetylmethylamine (AEI) inactivated vaccine no animal was positive for VIAA. Pinto and Garland¹⁰ later found VIAA positive cases also in animals revaccinated with AEI inactivated vaccine, but emphasized that the response in these animals was much weaker and only transient. Alonso *et al.*¹¹ were able to confirm this transient and weak response and also found, with binary ethylenimine (BEI) inactivated Al(OH)₃ and oil adjuvant vaccines, a response to VIAA only after revaccination. The weak response to VIAA is caused by the presence of this antigen in the inactivated virus suspension used for vaccine preparation. However VIAA antibodies induced by safely inactivated vaccines are only detected after revaccination and the presence of such antibodies after primovaccination still indicates recent virus replication in the animal.

Lucam *et al.* analysed in 1958 the FMD vaccination

Pan American Foot-and-Mouth Disease Center, Pan American Health Organization—WHO, Caixa Postal 569, 20001 Rio de Janeiro, Brasil. (Received 9 October 1989; revised 6 December 1989; accepted 6 December 1989)

campaigns in France¹² and suspected residual infectivity in some formaldehyde inactivated FMD vaccines, which had passed official innocuity tests.

Recently Beck and Strohmaier¹³ studied viruses from field outbreaks of FMD in Europe by determination of their nucleotide sequences. They found that most of these isolates were related to virus strains in (formaldehyde inactivated) vaccines. This led Strohmaier¹⁴ to the opinion that most of the FMD outbreaks in Europe in the last 20 years were 'homemade', i.e. were caused by vaccination. He made the recommendation that inactivation of FMD vaccine antigens should be changed from formaldehyde to first-order inactivants or that vaccination should be stopped altogether.

The FMD vaccine production regulations in several South American countries have for several years now permitted only the use of first-order inactivants. At least one European country has recently also adopted this position. Most of the FMD vaccine production laboratories in these countries apply the aziridine compound EI in the form of BEI. One group of laboratories uses diluted EI.

Inactivation with aziridines

The first report of a (bacterial) virus inactivation by ethylenimine, the basic aziridine substance, was published in 1955 by Raettig and Uecker¹⁵. Hurst in 1957¹⁶ was of the opinion that vaccines prepared with AEI as inactivant were antigenically superior to vaccines inactivated with formaldehyde and would guarantee inactivation of the virus. The antigenic superiority was later confirmed also for vaccines inactivated with BEI¹⁷⁻¹⁹. ICI patented the use of AEI for inactivation of microorganisms in 1959²⁰. The first report on the inactivation of FMD virus by AEI was published by Brown and Crick also in 1959²¹. This compound was subsequently used by a leading FMD vaccine production laboratory for many years in the preparation of inactivated antigens. However it did not come into general use because of its patent protection.

In 1961 Uecker²² reported the linearity of inactivation of bacterial viruses by ethylenimine derivatives and Graves and Arlinghaus described in 1967 the linearity of AEI inactivation of foot-and-mouth disease virus²³.

At ambient temperatures AEI is not stable and it therefore has to be kept at 4°C or preferably at -20°C. Fellows remarked in 1965²⁴, that AEI has a low boiling point and very little is left in a biologic preparation at reaction temperatures of 20°C or above. This observation probably made him use an inactivation temperature of 23°C instead of the usual 37°C.

The problem of the stability of AEI was perhaps the reason for introducing the double dosing regimen for inactivation at 37°C, i.e. the application of two doses for 24 h each as described by Pay *et al.*²⁵. This procedure is still being used with BEI by the same laboratory²⁶ although EI is much more stable²⁷. An extended incubation of the antigen at 37°C damages the antigen, as was shown in comparative inactivations of FMD virus at 26°C and at 37°C²⁸. This damage is not due to the inactivant but is probably caused by the action of proteolytic enzymes present in the virus suspension.

Other laboratories continued to work on FMD virus inactivation by aziridines and in the early 1970s reports on ethylethylenimine, EEI²⁹, ethylenimine, EI²⁷ and binary ethylenimine, BEI³⁰ were published. Both EEI as

well as EI are difficult to obtain in quantity. For this reason, as well as the ease of preparation and handling, BEI is now the preferred inactivating agent for FMD and other veterinary vaccines.

The viruses which have been reported as inactivated with BEI are given in Table 1. They belong to a variety of families of viruses with either RNA or DNA, which makes it very likely that most known viruses would be inactivated by an aziridine.

Inactivation with BEI

General considerations

Inactivation in vaccine preparation transforms an infectious antigen into a non-infectious one. This transformation step should therefore be done in a well identified intermediate area between the virus-containing and the virus-free area. Access to the intermediate area should be limited, and only be possible from the virus-containing side.

The antigen must be held in the intermediate area until completion of the necessary control tests (inactivation endpoint and innocuity). The facilities for holding of the inactivated antigen in this area, cold room or cooled storage tanks, should be able to accommodate a volume of at least 2-3 weeks production of virus in order to allow termination and if necessary a repeat of the control tests. Only after confirmation of the innocuity can the antigen be transferred to the virus-free area and be used for vaccine preparation.

The inactivation process should be done under slow agitation in two different vessels, with perhaps one quarter or a third of the time in the first vessel, and transfer of the virus suspension under inactivation in a closed system to the second vessel for the remainder of the time. This procedure is indicated in order to avoid pockets of the virus suspension into which the inactivant did not enter or reinfection from non-inactivated virus on the tank wall above the liquid level at the end of the inactivation period and after hydrolyzation of the inactivant.

The virus suspension should be checked to determine that it is at the desired temperature and is at a pH of ≈ 7.4 before the inactivant is added. It is also advisable to control the osmolality of the virus suspension, which for cell culture produced FMD virus is $\approx 320-340$ mOsm.

Table 1 Viruses inactivated with BEI

Virus	Nucleic acid	Ref.
1 African swine fever	DNA	31
2 Bluetongue	RNA	32
3 Bovine leukoemia	RNA	33
4 Bovine rhinotracheitis	DNA	34
5 Bovine rhinovirus	RNA	35
6 Bovine viral diarrhoea	RNA	36
7 Eastern equine encephalomyelitis	RNA	Unpublished
8 Foot-and-mouth disease	RNA	30
9 Newcastle disease	RNA	18
10 Porcine parvovirus	DNA	37
11 Pseudorabies	DNA	38
12 Rabies virus	RNA	39
13 Vesicular stomatitis	RNA	34

log₁₀ inactivation rate (min)

Figure 1

log₁₀ inactivation rate (min)

Figure 2

The result osmo measurements can reduce forms was: Figure down on the matio the p

Prep The becau ethyla distim in an active comp The NaOH 30-60 from: contr indica stock:

Inactivation of viral antigens: H. G. Bahnemann

log 10⁻³ (for the usually expressed titre in ml). The minimum endpoint for this volume therefore is log 10⁻⁴. For a successful inactivation the calculated endpoint has to be lower than the minimum endpoint. The difference between the calculated and the minimum endpoint, or the DIM (Difference of Inactivation endpoint to Minimum) value has to be positive. It is important to determine the DIM value for each inactivation process for an assessment of a successful inactivation.

After termination of the inactivation each virus suspension has to be tested for innocuity on cell cultures. This can be done by inoculation of at least two roller bottles or similar cell culture vessels and two subsequent blind passages at 48 h intervals. The cell cultures for this test have to be prepared in the virus-free area and the test should be done in the intermediate area. Only after determination of the inactivation endpoint and completion of the innocuity test can the virus suspension be considered to be properly inactivated.

Other applications of BEI

The major application of BEI will be in the preparation of inactivated antigen for vaccines. But there are other areas in which BEI can be used. Since BEI does not react with proteins it can be used for inactivation of adventitious viruses in biological preparations from animal or human tissues or fluids.

The BEI treatment of bovine serum used for cell cultures has been reported in 1976³⁴. The usefulness of this method was later confirmed in 1984 by Heuschele³⁵, who applied it over a 6-year period for the inactivation of adventitious bovine viral diarrhoea virus in calf serum used in primary or secondary animal cell cultures.

BEI can also be used for inactivation of viruses in enzyme preparations of animal origin. A commercial trypsin preparation was treated with BEI without any loss of activity (unpublished results). It is very likely that other biologicals, like Factor VIII, could be treated with BEI for inactivation of adventitious viruses.

Safety

Pure aziridines are highly toxic and have to be handled with special precautions and extreme care. This high toxicity is the reason for the 0.1 M preparation of BEI. At this molarity the BEI preparation contains only 0.5% EI and the vapour pressure at this concentration is low enough that for temperatures under 50°C no EI will escape into the atmosphere (the boiling point for EI is 57°C).

On the basis of experiments in laboratory animals, aziridines are considered to be carcinogenic substances. According to Derner and Ham⁴³, however, no cases of human cancer caused by EI have ever been reported. Fellowes⁴⁴ cites Hurst who states that an injection of 0.5 mg of AEI into rats did not produce any tumours during an observation period of 515 days.

The total annual production of FMD vaccine worldwide is probably between 700 and 800 million doses, of which nearly 500 million doses are produced in South America alone. Furthermore between 70% and 80% of all FMD vaccines are inactivated with BEI, which means that about 500 million doses of FMD vaccine are inactivated with BEI and applied in cattle annually. The majority of these cattle are revaccinated many times over the years. No increase in the incidence of cancer in cattle




Figure 3 Screen menu of the computer program for inactivation

after vaccination with BEI inactivated vaccines has ever been reported from any country. Neither is there any reason to expect any increase, as the residual BEI after inactivation is hydrolyzed with Na-thiosulphate.

Computer program

A computer program has been written to assist with the calculations needed for a controlled inactivation of viruses with BEI or any other first-order inactivant. The screen menu for the program is given in Figure 3. The program runs with DOS and is available in English (vine.exe) or in Spanish (vins.exe). Copies of either program can be obtained by sending a formatted double-density/double sided 5.25 in. diskette. Requests from South America should be sent to the Centro Panamericano de Febre Aftosa, Caixa Postal 589, 20001 Rio de Janeiro, Brasil. Requests from North America, Europe or other countries can be sent to the Pan American Health Organization, attention DIC/USA, 525-23rd Street, NW, Washington, DC 20037, USA.

References

1. Weselén, T., Lycke, E., Gard, S. and Ölin, G. Inactivation of poliomyelitis virus by formaldehyde. *Arch. Ges. Virus.* 1957, 1, 125-126.
2. Gard, S. and Lycke, E. Inactivation of poliovirus by formaldehyde. Analysis of inactivation curves. *Arch. Ges. Virus.* 1957, 1, 471-483.
3. Weselén, T. and Dinter, Z. The inactivation of foot-and-mouth disease virus by formalin. *Arch. Ges. Virus.* 1957, 1, 394-401.
4. Graves, J.H. Formaldehyde inactivation of foot-and-mouth disease virus as applied to vaccine preparation. *Am. J. Vet. Res.* 1963, 24, 1131-1135.
5. Barteling, S.J. and Wootmeyster, R. Formaldehyde inactivation of foot-and-mouth disease virus. Conditions for the preparation of safe vaccine. *Arch. Virus.* 1964, 88, 103-117.
6. Schultz, P., Righter, W.A., Timm, E.A., Taylor, A.R. and McLean, Jr. L.W. Partially inactivated poliomyelitis virus: Initiation of infection in tissue culture. *J. Immunol.* 1957, 78, 497-507.
7. Böfinger, M., Lycke, E., Möller, B. and Wrangé, G. Inactivation of poliomyelitis virus by formaldehyde. Incubation time in tissue culture of formalin treated virus. *Arch. Ges. Virus.* 1958, 8, 259-268.
8. Polietnick, J., Arlinghaus, R., Graves, J.H. and Cowan, K.M. Inhibition of cell-free foot-and-mouth disease virus ribonucleic acid synthesis by antibody. *Virology* 1967, 31, 600-615.
9. Alonso, A., de Mello, P.A., Gomes, I. and Rosenberg, F. The use of virus-infection-associated antigen (VIA) in the detection of cattle exposed to foot-and-mouth disease. *Bull. C.P.F.A.* 1973, 17-18, 17-34.
10. Pinto, A.A. and Garland, A.J.M. Immune response to virus-infection-associated (VIA) antigen in cattle repeatedly vaccinated with foot-and-mouth disease virus inactivated by formalin or acetyl ethylenimine. *J. Hyg. Camb.* 1979, 82, 41-50.

O
car

[1986/1986]

activation

nes has eve
is there any
al BEI after
late.

assist with
activation of
divant. The
gure 3. The
in English
s of either
formatted
Request
the Centro
589, 20001
h America,
o the Pan
DIC/USA,
7, USA.

activation of
1987, 7.

ormaldehyde.
7, 471-493
mouth disease

mouth disease
1963, 34

activation of
fection of pigs

and McLean,
n of infection

activation of
issue culture
53-255

JA. Inhibition
did synthesis

. F. The use
tion of cattle
17-18, 17-24
us-infection
inhalated with
or acetyl-

- 11 Alonso, A., Gomes, I. and Bahnmann, H.G. The induction of antibodies against VIAA in cattle vaccinated and revaccinated with inactivated FMD vaccine. *Bull. C.P.P.A.* 1988, 54, 33-34 in press
- 12 Lucem, F., Mackowiak, C. and Magat, A. La vaccination entiphouse en France. *Bull. O.J.E.* 1958, 48, 230-235
- 13 Beck, E. and Strohmair, K. Subtyping of European foot-and-mouth disease strains by nucleotide sequence determination. *J. Virol.* 1987, 61, 1821-1829
- 14 Strohmair, K. Die Maul- und Klauenseuche III. Erfahrungen und Folgerungen aus 20 Jahren Bekämpfung der Maul- und Klauenseuche. *Tierärztl. Umschau* 1987, 42, 417-422
- 15 Riedig, H. and Uecker, W. Inaktivierung von Bakteriophagen durch Äthyleniminderstoffe. *Naturwissenschaften* 1966, 42, 480
- 16 Hurst, A. Approaches to the chemotherapy of virus diseases. *J. Pharm. Pharmacol.* 1957, 8, 273-292
- 17 Girard, H.C., Beyrasmoglu, O., Erol, N. and Burgut, A. Inactivation of O, FMD virus by binary ethylenimine (BEI). *Bull. O.J.E.* 1977, 87, 201-217
- 18 Makarasin, P., Sinsuwonkiet, P. and Tanacharoonwich, P. Binary ethylenimine and foot-and-mouth disease vaccine production in Thailand. In: *Veterinary Viral Diseases: Their Significance in South-East Asia and the Western Pacific*, Academic Press, Australia, 1985, pp 308-309
- 19 Buonavoglia, C., Floretti, A., Tolla, M., Menna, F. and Paparella, V. A preliminary vaccine potency trial of a Newcastle disease virus inactivated with binary ethylenimine. *Vet. Rec. Comm.* 1988, 12, 185-187
- 20 ICZ Br. Pat. 608 949 and 608 950 (1959)
- 21 Brown, F. and Crick, J. Application of agar-gel diffusion analysis to a study of the antigenic structure of inactivated vaccines prepared from the virus of foot-and-mouth disease. *J. Immunol.* 1958, 82, 444-447
- 22 Uecker, W. Inaktivierung von Mikroorganismen durch Äthylierende Verbindungen. I. Untersuchungen zur Kinetik der Inaktivierung von Bakteriophagen. *Zentralbl. Bakteriol. Hyg. Abt. I. Orig.* 1961, 181, 518-531
- 23 Graves, J.H. and Arlinghaus, R.B. Acetylthymine in the preparation of inactivated foot-and-mouth disease vaccines. *Proc. 71st Ann. Meeting US Livest. Sanit. Assoc.* 1987, pp 396-403
- 24 Fellows, O.H. Comparison of the inactivation and antigenicity of foot-and-mouth disease virus by acetylthymine and by combined effect of ultraviolet light and β -propiolactone. *J. Immunol.* 1963, 98, 1100-1105
- 25 Pay, T.W.F. et al. Some observations on foot-and-mouth disease virus inactivation with acetylthymine. *Eur. Foot-and-Mouth Disease Commission, Meeting Technical Committee, Tübingen, 1977, FAO, Rome*
- 26 Pay, T.W.F., Telling, R.C. and Thoms, A.C. Inactivation of FMD virus with aziridines. *Eur. Foot-and-Mouth Disease Commission, Meeting Technical Committee, Tübingen, 1981, FAO, Rome*
- 27 Bahnmann, H.G. The inactivation of foot-and-mouth disease virus by ethylenimine and propylenimine. *Zentralbl. Vet. Med. B* 1973, 28, 336-350
- 28 Bahnmann, H.G., Mesquita, J., Astudillo, V. and Dora, F. The production and application of an oil adjuvant vaccine against

Inactivation of viral antigens: H. G. Bahnmann

- foot-and-mouth disease in cattle. In: *Modern Approaches to Animal Cell Technology* (Eds Epier, R.E. and Griffiths, J.B.) Butterworth & Co., UK, 1987
- 29 Bauer, K. Die Inaktivierung des Maul- und Klauenseuche (MKS) Virus durch Äthylenimine und die Eignung des inaktivierten Virus zur Impfstoffherstellung. *Zentralbl. Bakteriol. Hyg. Abt. I Orig.* 1970, 213, 285-297
- 30 Bahnmann, H.G. Binary ethylenimine as an inactivant for foot-and-mouth disease virus and its application for vaccine production. *Arch. Virol.* 1975, 47, 47-56
- 31 Schloer, G.M. Kinetics of inactivation of African swine fever antigen with binary ethylenimine. *Proc. US Anim. Health Assoc.* 1982, 86, 233-238
- 32 Barber, T.L. and Jochim, M.J. Inactivation of bluetongue virus with binary ethylenimine. *Ann. Meet. Am. Soc. Microbiol.* 1978, p 339
- 33 Miller, J.M. and Van der Maaten, M.J. Evaluation of an inactivated bovine leukemia virus preparation as an immunogen in cattle. *Ann. Rech. Vet.* 1978, 8, 871-877
- 34 Bahnmann, H.G. Inactivation of viruses in serum with binary ethylenimine. *J. Clin. Microbiol.* 1975, 3, 208-210
- 35 Crouch, C.F. Vaccination against enteric rota and coronaviruses in cattle and pigs: enhancement of isogenic immunity. *Vaccine* 1985, 3, 284-291
- 36 Hauschke, W.P. Inactivation of adventitious viruses in serum. *Art. Science* 1984, 3, 2
- 37 Wraith, A.E., Wells, D.E., Cartwright, S.F. and Frerichs, G.N. An inactivated oil emulsion vaccine for the prevention of porcine parvovirus-induced reproductive failure. *Res. Vet. Sci.* 1984, 36, 136-143
- 38 Sun, L.L., Gustafson, D.P. and Scheraga, G. Comparison of pseudorabies virus inactivated by bromo-ethylenimine. "Co irradiation and acridine dye in immune assay systems. *J. Clin. Microbiol.* 1978, 8, 804-811
- 39 Lough, G.P. and Nebel, A.E. Rabies virus inactivation by binary ethylenimine: new method for inactivated vaccine production. *J. Clin. Microbiol.* 1980, 11, 120-122
- 40 Lycke, E. Studies of the inactivation of poliovirus by formaldehyde. The effect upon the rate of inactivation of temperature, pH and concentration of formaldehyde. *Arch. Ges. Virol.* 1958, 8, 257-264
- 41 Epstein, J., Rosenthal, R.W. and Ess, R.J. Use of γ -44-nitrobenzylpyridine as analytical reagent for ethylenimines and other alkylating agents. *Anal. Chem.* 1953, 27, 1435-1439
- 42 Collong, F., Farkas, L., Perenyi, T., Fazekas, A., Szalai, F., Zsitvai, K. and Erdelyi, K. Effect of ethylenimine and formaldehyde inactivation on the immunogenicity of foot-and-mouth disease virus. *XVth Conf. O.J.E. Commission on Foot-and-Mouth Disease, Paris 10-14 October 1978*
- 43 Pseud, J., Dubourget, P., Detraz, N. and Lombard, M. Inactivation of Foot-and-Mouth Disease Virus by Ethylenimine - Kinetics and Control. *Int. Foot-and-Mouth Disease Symp. Ankara, Turkey, June 1988 FMD Institute Ankara*
- 44 Darmer, G.C. and Ham, G.E. *Ethylenimine and other Aziridines*. Academic Press, New York, 1968, p 400

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ BLACK BORDERS

☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.